arXiv:1107.4033v1 [math.CA] 20 Jul 2011

ON THE CO-ORDINATED CONVEX FUNCTIONS

M. EMIN ÖZDEMIR■, ÇETIN YILDIZ■♠, AND AHMET OCAK AKDEMIR★

Abstract. In this paper we established new integral inequalities which are more general results for co-ordinated convex functions on the co-ordinates by using some classical inequalities.

1. INTRODUCTION

Let $f : I \subseteq \mathbb{R} \to \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a < b$. The following double inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}$$

is well known in the literature as Hadamard's inequality. Both inequalities hold in the reversed direction if f is concave.

In [4], Dragomir defined convex functions on the co-ordinates as following;

Definition 1. *Let us consider the bidimensional interval* $\Delta = [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$, $c < d$. *A function* $f : \Delta \to \mathbb{R}$ *will be called convex on the co-ordinates if the partial mappings* $f_y : [a,b] \to \mathbb{R}$, $f_y(u) = f(u,y)$ *and* $f_x : [c,d] \to \mathbb{R}$, $f_x(v) = f(x,v)$ *are convex where defined for all* $y \in [c,d]$ *and* $x \in [a,b]$. *Recall that the mapping* $f : \Delta \to \mathbb{R}$ *is convex on* Δ *if the following inequality holds,*

$$f(\lambda x + (1-\lambda)z, \lambda y + (1-\lambda)w) \leq \lambda f(x,y) + (1-\lambda)f(z,w)$$

for all $(x,y),(z,w) \in \Delta$ *and* $\lambda \in [0,1]$.

In [4], Dragomir established the following inequalities of Hadamard's type for co-ordinated convex functions on a rectangle from the plane $\mathbb{R}^2$.

Date: April, 2011.

2000 *Mathematics Subject Classification.* Primary 26D15.

Key words and phrases. Hadamard's inequality, co-ordinates, convexity, Hölder's inequality, Power mean inequality.

♠corresponding author.

Theorem 1. *Suppose that* $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ *is convex on the co-ordinates on* Δ*. Then one has the inequalities;*

$$
\begin{aligned}
& f(\frac{a+b}{2},\frac{c+d}{2}) \\
\leq\ & \frac{1}{2}\left[\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx+\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy\right] \\
\leq\ & \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy \\
\leq\ & \frac{1}{4}\left[\frac{1}{(b-a)}\int_a^b f(x,c)dx+\frac{1}{(b-a)}\int_a^b f(x,d)dx\right. \\
& \left.+\frac{1}{(d-c)}\int_c^d f(a,y)dy+\frac{1}{(d-c)}\int_c^d f(b,y)dy\right] \\
\leq\ & \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}.
\end{aligned}
\tag{1.1}
$$

The above inequalities are sharp.

In [1], Bakula and Pečarić established several Jensen type inequalities for co-ordinated convex functions and in [5], Hwang *et al.* gave a mapping F, discussed some properties of this mapping and proved some Hadamard-type inequalities for Lipschizian mapping in two variables. In [2], Özdemir *et al.* established new Hadamard-type inequalities for co-ordinated $m-$convex and $(\alpha,m)-$convex functions. On all of these, in [3], the authors proved some Hadamard-type inequalities for co-ordinated convex functions as followings;

Theorem 2. *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta := [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$ *and* $c < d$*. If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ *is a convex function on the co-ordinates on* Δ*, then one has the inequalities:*

$$
\begin{aligned}
& \left|\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right. \\
& \left.\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy - A\right| \\
\leq\ & \frac{(b-a)(d-c)}{16}\left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}\right|(a,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(a,d)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(b,c)+\left|\frac{\partial^2 f}{\partial t\partial s}\right|(b,d)}{4}\right)
\end{aligned}
\tag{1.2}
$$

where

$$
A=\frac{1}{2}\left[\frac{1}{(b-a)}\int_a^b\left[f(x,c)+f(x,d)\right]dx+\frac{1}{(d-c)}\int_c^d\left[f(a,y)dy+f(b,y)\right]dy\right].
$$

Theorem 3. *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta := [a,b] \times [c,d]$ *in* $\mathbb{R}^2$ *with* $a < b$ *and* $c < d$*. If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$*,* $q > 1$*, is a convex function on* *the co-ordinates on* Δ, *then one has the inequalities:*

$$\begin{aligned} (1.3)\quad & \left| \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \right. \\ & \left. \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)dxdy - A \right| \\ \leq \quad & \frac{(b-a)(d-c)}{4\,(p+1)^{\frac{2}{p}}} \left(\frac{\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (a,c) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (a,d) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (b,c) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (b,d)}{4} \right)^{\frac{1}{q}} \end{aligned}$$

where

$$A = \frac{1}{2}\left[\frac{1}{(b-a)} \int_a^b \left[f(x,c)+f(x,d)\right] dx + \frac{1}{(d-c)} \int_c^d \left[f(a,y)dy + f(b,y)\right] dy \right]$$

and $\frac{1}{p}+\frac{1}{q}=1$.

Theorem 4. *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta := [a,b]\times[c,d]$ *in* $\mathbb{R}^2$ *with* $a<b$ *and* $c<d$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$, $q \geq 1$, *is a convex function on the co-ordinates on* Δ, *then one has the inequalities:*

$$\begin{aligned} (1.4)\quad & \left| \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \right. \\ & \left. \frac{1}{(b-a)(d-c)} \int_a^b \int_c^d f(x,y)dxdy - A \right| \\ \leq \quad & \frac{(b-a)(d-c)}{16} \left(\frac{\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (a,c) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (a,d) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (b,c) + \left|\frac{\partial^2 f}{\partial t \partial s}\right|^q (b,d)}{4} \right)^{\frac{1}{q}} \end{aligned}$$

where

$$A = \frac{1}{2}\left[\frac{1}{(b-a)} \int_a^b \left[f(x,c)+f(x,d)\right] dx + \frac{1}{(d-c)} \int_c^d \left[f(a,y)dy + f(b,y)\right] dy \right].$$

In [6], authors proved following inequalities for co-ordinated convex functions.

Theorem 5. *Let* $f : \Delta = [a,b]\times[c,d] \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta = [a,b]\times[c,d]$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|$ *is a convex function on the co-ordinates on* Δ, *then the following inequality holds;*

$$\begin{aligned} (1.5)\quad & \left| f\left(\frac{a+b}{2}, \frac{c+d}{2}\right) \right. \\ & -\frac{1}{(d-c)} \int_c^d f\left(\frac{a+b}{2}, y\right) dy - \frac{1}{(b-a)} \int_a^b f\left(x, \frac{c+d}{2}\right) dx \\ & \left. +\frac{1}{(b-a)\,(d-c)} \int_a^b \int_c^d f\,(x,y)\,dydx \right| \\ \leq \quad & \frac{(b-a)\,(d-c)}{64} \left[\left|\frac{\partial^2 f}{\partial t \partial s}\,(a,c)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\,(b,c)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\,(a,d)\right| + \left|\frac{\partial^2 f}{\partial t \partial s}\,(b,d)\right| \right]. \end{aligned}$$

Theorem 6. *Let* $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta = [a,b] \times [c,d]$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$, $q > 1$, *is a convex function on the co-ordinates on* Δ, *then the following inequality holds;*

$$\begin{aligned}
&\left| f\left(\frac{a+b}{2},\frac{c+d}{2}\right) + \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx \right. \\
&\left. -\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy - \frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \right| \\
\leq\ & \frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}} \\
&\times \left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4} \right)^{\frac{1}{q}}.
\end{aligned} \tag{1.6}$$

Theorem 7. *Let* $f : \Delta = [a,b] \times [c,d] \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta = [a,b] \times [c,d]$. *If* $\left|\frac{\partial^2 f}{\partial t \partial s}\right|^q$, $q \geq 1$, *is a convex function on the co-ordinates on* Δ, *then the following inequality holds;*

$$\begin{aligned}
&\left| f\left(\frac{a+b}{2},\frac{c+d}{2}\right) + \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx \right. \\
&\left. -\frac{1}{(d-c)}\int_c^d f\left(\frac{a+b}{2},y\right)dy - \frac{1}{(b-a)}\int_a^b f\left(x,\frac{c+d}{2}\right)dx \right| \\
\leq\ & \frac{(b-a)(d-c)}{16} \\
&\times \left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right|^q + \left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|^q}{4} \right)^{\frac{1}{q}}.
\end{aligned} \tag{1.7}$$

In [7], Özdemir *et al.* proved the following Theorem which is involving an inequality of Simpson's type;

Theorem 8. *Let* $f : \Delta \subset \mathbb{R}^2 \to \mathbb{R}$ *be a partial differentiable mapping on* $\Delta = [a,b] \times [c,d]$. *If* $\frac{\partial^2 f}{\partial t \partial s}$ *is a convex function on the co-ordinates on* Δ, *then the following inequality holds:*

$$\begin{aligned}
&\left| \frac{f\left(a,\frac{c+d}{2}\right) + f\left(b,\frac{c+d}{2}\right) + 4f\left(\frac{a+b}{2},\frac{c+d}{2}\right) + f\left(\frac{a+b}{2},c\right) + f\left(\frac{a+b}{2},d\right)}{9} \right. \\
&\left. + \frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36} + \frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx - A \right| \\
\leq\ & \frac{25(b-a)(d-c)}{72} \\
&\times \left(\frac{\left|\frac{\partial^2 f}{\partial t\partial s}(a,c)\right| + \left|\frac{\partial^2 f}{\partial t\partial s}(a,d)\right| + \left|\frac{\partial^2 f}{\partial t\partial s}(b,c)\right| + \left|\frac{\partial^2 f}{\partial t\partial s}(b,d)\right|}{72} \right)
\end{aligned} \tag{1.8}$$

where

$$\begin{aligned} A &= \frac{1}{6(b-a)}\int_a^b\left[f(x,c)+4f\left(x,\frac{c+d}{2}\right)+f(x,d)\right]dx \\ &\quad+\frac{1}{6(d-c)}\int_c^d\left[f(a,y)+4f\left(\frac{a+b}{2},y\right)+f(b,y)\right]dy. \end{aligned}$$

The main purpose of this paper is to establish a new lemma which give more general results and different type inequalities for special values of λ and to prove several inequalities.

2. MAIN RESULTS

In order to prove our main theorems we need the following lemma:

Lemma 1. *Let $f:\Delta\subset\mathbb{R}^2\to\mathbb{R}$ be a differentiable function on Δ where $a<b$, $c<d$ and $\lambda\in[0,1]$, if $\frac{\partial^2 f}{\partial x\partial y}\in L_1(\Delta)$, then the following equality holds:*

$$\begin{aligned} &(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \\ &+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right] \\ &-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx \\ &-\lambda\frac{1}{2(b-a)}\int_a^b[f(x,d)+f(x,c)]dx-\lambda\frac{1}{2(d-c)}\int_c^d[f(a,y)+f(b,y)]dy \\ &+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy \\ =\ &\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d K(x)M(y)\frac{\partial^2 f}{\partial x\partial y}(x,y)dydx \end{aligned}$$

where

$$K(x)=\begin{cases} x-\left(a+\lambda\frac{b-a}{2}\right), & x\in\left[a,\frac{a+b}{2}\right] \\ x-\left(b-\lambda\frac{b-a}{2}\right), & x\in\left[\frac{a+b}{2},b\right] \end{cases}$$

and

$$M(y)=\begin{cases} y-\left(c+\lambda\frac{d-c}{2}\right), & y\in\left[c,\frac{c+d}{2}\right] \\ y-\left(d-\lambda\frac{d-c}{2}\right), & y\in\left[\frac{c+d}{2},d\right] \end{cases}.$$

Proof. Integration by parts, we get

$$\begin{aligned}
&\int_a^b\int_c^d K(x)M(y)\frac{\partial^2 f}{\partial x\partial y}(x,y)dydx\\
=&\int_a^b K(x)\left[\int_c^{\frac{c+d}{2}}\left(y-\left(c+\lambda\frac{d-c}{2}\right)\right)\frac{\partial^2 f}{\partial x\partial y}(x,y)dy\right.\\
&\left.+\int_{\frac{c+d}{2}}^d\left(y-\left(d-\lambda\frac{d-c}{2}\right)\right)\frac{\partial^2 f}{\partial x\partial y}(x,y)dy\right]dx\\
=&\int_a^b K(x)\left[\left(y-\left(c+\lambda\frac{d-c}{2}\right)\right)\frac{\partial f}{\partial x}(x,y)\Big|_c^{\frac{c+d}{2}}-\int_c^{\frac{c+d}{2}}\frac{\partial f}{\partial x}(x,y)dy\right.\\
&\left.+\left(y-\left(d-\lambda\frac{d-c}{2}\right)\right)\frac{\partial f}{\partial x}(x,y)\Big|_{\frac{c+d}{2}}^d-\int_{\frac{c+d}{2}}^d\frac{\partial f}{\partial x}(x,y)dy\right]dx\\
=&\int_a^b K(x)\left[(1-\lambda)(d-c)\frac{\partial f}{\partial x}\left(x,\frac{c+d}{2}\right)\right.\\
&\left.+\left(\lambda\frac{d-c}{2}\right)\left(\frac{\partial f}{\partial x}(x,c)+\frac{\partial f}{\partial x}(x,d)\right)-\int_c^d\frac{\partial f}{\partial x}(x,y)dy\right]dx.
\end{aligned}$$

Integrating by parts again, we obtain

$$\begin{aligned}
&\int_a^b\int_c^d K(x)M(y)\frac{\partial^2 f}{\partial x\partial y}(x,y)dydx\\
=&(1-\lambda)^2(b-a)(d-c)\left[f\left(\frac{a+b}{2},\frac{c+d}{2}\right)\right]\\
&+\lambda^2(b-a)(d-c)\left[\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right]\\
&+\frac{\lambda(1-\lambda)(b-a)(d-c)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)\right.\\
&\left.+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]-(1-\lambda)(b-a)\int_c^d f(\frac{a+b}{2},y)dy\\
&-(1-\lambda)(d-c)\int_a^b f(x,\frac{c+d}{2})dx-\lambda\frac{(d-c)}{2}\int_a^b[f(x,d)+f(x,c)]dx\\
&-\lambda\frac{(b-a)}{2}\int_c^d[f(a,y)+f(b,y)]dy+\int_a^b\int_c^d f(x,y)dxdy.
\end{aligned}$$

Dividing both sides of the above equality by $(b-a)(d-c)$, we have the required result. □

Theorem 9. *Let $f:\Delta=[a,b]\times[c,d]\to\mathbb{R}$ be a differentiable function on Δ. If $\left|\frac{\partial^2 f}{\partial x\partial y}\right|$ is convex function on the co-ordinates on Δ, then one has the inequality:*

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\;&\frac{(b-a)(d-c)}{16}\left[2\lambda^2-2\lambda+1\right]^2\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|(b,d)}{4}\right).
\end{aligned}$$

Proof. From Lemma 1 and property of the modulus, we can write

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\;&\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d\left|K(x)M(y)\right|\left|\frac{\partial^2 f}{\partial x\partial y}\right|(x,y)dydx.
\end{aligned}$$

By using the change of variables $y = sd + (1-s)\,c$, $(d-c)\,ds = dy$, we obatin

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right) + \lambda^2 \frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right] \\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx \\
&-\lambda\frac{1}{2(b-a)}\int_a^b [f(x,d)+f(x,c)]\,dx-\lambda\frac{1}{2(d-c)}\int_c^d [f(a,y)+f(b,y)]\,dy \\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg| \\
\leq\ &\frac{d-c}{b-a}\int_a^b |K(x)|\left\{\int_0^{\frac{\lambda}{2}}\left(\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,sd+(1-s)\,c)\right| ds\right. \\
&+\int_{\frac{\lambda}{2}}^{\frac{1}{2}}\left(s-\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,sd+(1-s)\,c)\right| ds \\
&+\int_{\frac{1}{2}}^{1-\frac{\lambda}{2}}\left(1-\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,sd+(1-s)\,c)\right| ds \\
&\left.+\int_{1-\frac{\lambda}{2}}^{1}\left(s-1+\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,sd+(1-s)\,c)\right| ds\right\} dx.
\end{aligned}$$

Since $\left|\frac{\partial^2 f}{\partial x\partial y}\right|$ is convex function on the co-ordinates on Δ, we have

$$\begin{aligned}
&\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d |K(x)M(y)|\left|\frac{\partial^2 f}{\partial x\partial y}\right|(x,y)dydx \\
\leq\ &\frac{d-c}{b-a}\int_a^b |K(x)|\left\{\int_0^{\frac{\lambda}{2}} s\left(\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,d)\right| ds\right. \\
&+\int_0^{\frac{\lambda}{2}}(1-s)\left(\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,c)\right| ds \\
&+\int_{\frac{\lambda}{2}}^{\frac{1}{2}} s\left(s-\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,d)\right| ds+\int_{\frac{\lambda}{2}}^{\frac{1}{2}}(1-s)\left(s-\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,c)\right| ds \\
&+\int_{\frac{1}{2}}^{1-\frac{\lambda}{2}} s\left(1-\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,d)\right| ds \\
&+\int_{\frac{1}{2}}^{1-\frac{\lambda}{2}}(1-s)\left(1-\frac{\lambda}{2}-s\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,c)\right| ds \\
&+\int_{1-\frac{\lambda}{2}}^{1} s\left(s-1+\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,d)\right| ds \\
&\left.+\int_{1-\frac{\lambda}{2}}^{1}(1-s)\left(s-1+\frac{\lambda}{2}\right)\left|\frac{\partial^2 f}{\partial x\partial y}(x,c)\right| ds\right\} dx.
\end{aligned}$$

By calculating the above integrals, we obtain

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\ &\frac{d-c}{b-a}\int_a^b|K(x)|\left\{\left[\frac{2\lambda^2-2\lambda+1}{8}\right]\left(\left|\frac{\partial^2 f}{\partial x\partial y}(x,c)\right|+\left|\frac{\partial^2 f}{\partial x\partial y}(x,d)\right|\right)\right\}.
\end{aligned}$$

By a similar argument for other integrals, by using the change of variable $x = tb+(1-t)a$, $(b-a)dt = dx$ and convexity of $\left|\frac{\partial^2 f}{\partial x\partial y}(x,y)\right|$ on the co-ordinates on Δ, we deduce the result. Which completes the proof. □

Remark 1. *If we choose* $\lambda = 1$ *in Theorem 5, we have the inequality (1.2).*

Remark 2. *If we choose* $\lambda = 0$ *in Theorem 5, we have the inequality (1.5).*

Remark 3. *If we choose* $\lambda = \frac{1}{3}$ *in Theorem 5, we have the inequality (1.8).*

Theorem 10. *Let* $f : \Delta = [a,b]\times[c,d] \to \mathbb{R}$ *be a differentiable function on* Δ. *If* $\left|\frac{\partial^2 f}{\partial x\partial y}\right|^{\frac{p}{p-1}}$ *is convex function on the co-ordinates on* Δ, *then one has the inequality:*

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\ &\frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left[2\lambda^2-2\lambda+1\right]^2\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}
\end{aligned}$$

for $q > 1$, *where* $q = \frac{p}{p-1}$.

Proof. Let $p>1$. From Lemma 1 and using the Hölder inequality for double integrals, we can write

$$\begin{aligned}
&\left|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right.\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b[f(x,d)+f(x,c)]\,dx-\lambda\frac{1}{2(d-c)}\int_c^d[f(a,y)+f(b,y)]\,dy\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right|\\
\leq\ &\frac{1}{(b-a)(d-c)}\left(\int_a^b\int_c^d|K(x)M(y)|^p\,dydx\right)^{\frac{1}{p}}\left(\int_a^b\int_c^d\left|\frac{\partial^2 f}{\partial x\partial y}(x,y)\right|^q dydx\right)^{\frac{1}{q}}.
\end{aligned}$$

Since $\left|\frac{\partial^2 f}{\partial x\partial y}\right|^{\frac{p}{p-1}}$ is convex function on the co-ordinates on Δ, by taking into account the change of variable $x=tb+(1-t)a$, $(b-a)dt=dt$ and $y=sd+(1-s)c$, $(d-c)ds=dy$, we have

$$\left|\frac{\partial^2 f}{\partial x\partial y}(tb+(1-t)a,y)\right|^q\leq t\left|\frac{\partial^2 f}{\partial x\partial y}(b,y)\right|^q+(1-t)\left|\frac{\partial^2 f}{\partial x\partial y}(a,y)\right|^q$$

and

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial x\partial y}(tb+(1-t)a,sd+(1-s)c)\right|^q\\
\leq\ & ts\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)+t(1-s)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)\\
&+s(1-t)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+(1-t)(1-s)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)
\end{aligned}$$

thus, we obtain

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b[f(x,d)+f(x,c)]\,dx-\lambda\frac{1}{2(d-c)}\int_c^d[f(a,y)+f(b,y)]\,dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\ &\frac{(b-a)(d-c)}{4(p+1)^{\frac{2}{p}}}\left[2\lambda^2-2\lambda+1\right]^2\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}.
\end{aligned}$$

Which completes the proof. □

Remark 4. *Under the assumptions of Theorem 6, if we choose* $\lambda=1$, *we have the inequality (1.3).*

Remark 5. *Under the assumptions of Theorem 6, if we choose* $\lambda=0$, *we have the inequality (1.6).*

Corollary 1. *Under the assumptions of Theorem 6, if we choose* $\lambda=\frac{1}{3}$, *we have the inequality*

$$\begin{aligned}
&\Bigg|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9}\\
&+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\Bigg|\\
\leq\ &\frac{25(b-a)(d-c)}{324(p+1)^{\frac{2}{p}}}\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}
\end{aligned}$$

where

$$\begin{aligned}
A\ =\ &\frac{1}{6(b-a)}\int_a^b\left[f(x,c)+4f\left(x,\frac{c+d}{2}\right)+f(x,d)\right]dx\\
&+\frac{1}{6(d-c)}\int_c^d\left[f(a,y)+4f\left(\frac{a+b}{2},y\right)+f(b,y)\right]dy.
\end{aligned}$$

Corollary 2. *In Corollary 1, since* $\frac{1}{4} < \frac{1}{(p+1)^{\frac{2}{p}}} < 1$*, for* $p > 1$*, we have the following inequality;*

$$\begin{aligned}
&\left| \frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9} \right.\\
&\left. +\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx - A\right| \\
\leq\ & \frac{25(b-a)(d-c)}{324} \\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}.
\end{aligned}$$

Theorem 11. *Let* $f : \Delta = [a,b]\times[c,d] \to \mathbb{R}$ *be a differentiable function on* Δ*. If* $\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q$ *is convex function on the co-ordinates on* Δ *and* $q \geq 1$*, then one has the inequality:*

$$\begin{aligned}
&\left|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\right.\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b[f(x,d)+f(x,c)]\,dx-\lambda\frac{1}{2(d-c)}\int_c^d[f(a,y)+f(b,y)]\,dy\\
&\left.+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\right|\\
\leq\ & \frac{(b-a)(d-c)}{16}\left[2\lambda^2-2\lambda+1\right]^2\\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}.
\end{aligned}$$

Proof. From Lemma 1 and using the well-known Power-mean inequality, we can write

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4}\\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right]\\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx\\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy\\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg|\\
\leq\;&\frac{1}{(b-a)(d-c)}\left(\int_a^b\int_c^d\left|K(x)M(y)\right|dydx\right)^{1-\frac{1}{q}}\\
&\times\left(\int_a^b\int_c^d\left|K(x)M(y)\right|\left|\frac{\partial^2 f}{\partial x\partial y}(x,y)\right|^q dydx\right)^{\frac{1}{q}}.
\end{aligned}$$

Since $\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q$ is convex function on the co-ordinates on Δ, by taking into account the change of variable $x=tb+(1-t)a$, $(b-a)dt=dt$ and $y=sd+(1-s)c$, $(d-c)ds=dy$, we have

$$\left|\frac{\partial^2 f}{\partial x\partial y}(tb+(1-t)a,y)\right|^q\leq t\left|\frac{\partial^2 f}{\partial x\partial y}(b,y)\right|^q+(1-t)\left|\frac{\partial^2 f}{\partial x\partial y}(a,y)\right|^q$$

and

$$\begin{aligned}
&\left|\frac{\partial^2 f}{\partial x\partial y}(tb+(1-t)a,sd+(1-s)c)\right|^q\\
\leq\;&ts\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)+t(1-s)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)\\
&+s(1-t)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+(1-t)(1-s)\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)
\end{aligned}$$

hence, it follows that

$$\begin{aligned}
&\Bigg|(1-\lambda)^2 f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+\lambda^2\frac{f(a,c)+f(a,d)+f(b,c)+f(b,d)}{4} \\
&+\frac{\lambda(1-\lambda)}{2}\left[f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)+f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)\right] \\
&-(1-\lambda)\frac{1}{d-c}\int_c^d f(\frac{a+b}{2},y)dy-(1-\lambda)\frac{1}{b-a}\int_a^b f(x,\frac{c+d}{2})dx \\
&-\lambda\frac{1}{2(b-a)}\int_a^b\left[f(x,d)+f(x,c)\right]dx-\lambda\frac{1}{2(d-c)}\int_c^d\left[f(a,y)+f(b,y)\right]dy \\
&+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)dxdy\Bigg| \\
\leq\ & \frac{(b-a)(d-c)}{16}\left[2\lambda^2-2\lambda+1\right]^2 \\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}.
\end{aligned}$$

Which completes the proof. □

Remark 6. *Under the assumptions of Theorem 7, if we choose* $\lambda=1$*, we have the inequality (1.4).*

Remark 7. *Under the assumptions of Theorem 7, if we choose* $\lambda=0$*, we have the inequality (1.7).*

Corollary 3. *Under the assumptions of Theorem 7, if we choose* $\lambda=\frac{1}{3}$*, we have*

$$\begin{aligned}
&\Bigg|\frac{f\left(a,\frac{c+d}{2}\right)+f\left(b,\frac{c+d}{2}\right)+4f\left(\frac{a+b}{2},\frac{c+d}{2}\right)+f\left(\frac{a+b}{2},c\right)+f\left(\frac{a+b}{2},d\right)}{9} \\
&+\frac{f(a,c)+f(b,c)+f(a,d)+f(b,d)}{36}+\frac{1}{(b-a)(d-c)}\int_a^b\int_c^d f(x,y)\,dydx-A\Bigg| \\
\leq\ & \frac{25(b-a)(d-c)}{36^2} \\
&\times\left(\frac{\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(a,d)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,c)+\left|\frac{\partial^2 f}{\partial x\partial y}\right|^q(b,d)}{4}\right)^{\frac{1}{q}}.
\end{aligned}$$

where

$$\begin{aligned}
A &= \frac{1}{6(b-a)}\int_a^b\left[f(x,c)+4f\left(x,\frac{c+d}{2}\right)+f(x,d)\right]dx \\
&\quad+\frac{1}{6(d-c)}\int_c^d\left[f(a,y)+4f\left(\frac{a+b}{2},y\right)+f(b,y)\right]dy.
\end{aligned}$$

References

[1] M.K. Bakula and J. Pečarić, On the Jensen's inequality for convex functions on the co-ordinates in a rectangle from the plane, *Taiwanese Journal of Math.*, 5, 2006, 1271-1292.

[2] M.E. Özdemir, E. Set, M.Z. Sarıkaya, Some new Hadamard's type inequalities for co-ordinated $m-$convex and $(\alpha, m)-$convex functions, Accepted.

[3] M.Z. Sarıkaya, E. Set, M. Emin Özdemir and S.S. Dragomir, New some Hadamard's type inequalities for co-ordinated convex functions, Accepted.

[4] S.S. Dragomir, On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane, *Taiwanese Journal of Math.*, 5, 2001, 775-788.

[5] D. Y. Hwang, K. L. Tseng and G. S. Yang, Some Hadamard's inequalities for co-ordinated convex functions in a rectangle from the plane, *Taiwanese Journal of Mathematics,* 11 (2007), 63-73.

[6] M.E. Özdemir, H. Kavurmacı, A.O. Akdemir and M. Avcı, Inequalities for convex and $s-$convex functions on $\Delta = [a, b] \times [c, d]$, Submitted.

[7] M.E. Özdemir, A.O. Akdemir and H. Kavurmacı, On the Simpson's inequality for co-ordinated convex functions, Submitted.

■Atatürk University, K.K. Education Faculty, Department of Mathematics, 25240, Campus, Erzurum, Turkey
E-mail address: `emos@atauni.edu.tr`

E-mail address: `yildizc@atauni.edu.tr`

★Ağri İbrahim Çeçen University, Faculty of Science and Letters, Department of Mathematics, 04100, Ağri, Turkey
E-mail address: `ahmetakdemir@agri.edu.tr`